

Target : Pre-Medical

Physics



28 years
of Excellence



0744-5156100 | www.allen.ac.in



Scanned with
CamScanner

*Contents***PHYSICS
RACES**

S.No.	Topic Name	Page No.
1	Units, Dimensions & Measurement	01
2	Friction	03
3	Work Power and Energy # 01	05
4	Work Power and Energy # 02	07
5	Circular motion	09
6	Collision and Centre of Mass	11
7	Rotational Motion # 01	13
8	Rotational Motion # 02	15
9	Rotational Motion # 03	17
10	Rotational Motion # 04	19
11	Gravitation	21
12	Properties of Matter & Fluid Mechanics (Elasticity) # 01	24
13	Properties of Matter & Fluid Mechanics (Fluid Dynamics + Viscosity) # 02	26
14	Properties of Matter & Fluid Mechanics (Hydrostatics) # 03	28
15	Properties of Matter & Fluid Mechanics (Surface Tension) # 04	30
16	Thermal Physics (Temperature Scales & Thermal Expansion) # 01	32
17	Thermal Physics (Calorimetry) # 02	34
18	Thermal Physics (Mode of Heat Transfer) # 03	36
19	Thermal Physics (Kinetic theory of gasses) # 04	39
20	Thermal Physics (Thermodynamic process) # 05	41
21	Oscillations (Kinematics of SHM) # 01	44
22	Oscillations (Energy & Spring Pendulum) # 02	46
23	Oscillations (Simple pendulum and types of SHM) # 03	48
24	Wave Motion & Doppler's Effect (Fundamental) #01	50
25	Wave Motion & Doppler's Effect (Superposition of waves interference, beats) # .02	53
26	Wave Motion & Doppler's Effect (Stationary waves & doppler effect, beats) # 03	55

Units, Dimensions & Measurements

- Which one of the following is not a fundamental quantity ?
(1) Temperature (2) Electric current
(3) Pressure (4) Length
- Which unit is used for measuring nuclear area of cross-section ?
(1) mm^2 (2) Fermi (3) Barn (4) Curie
- Which of the following is not correct representation of a unit ?
(1) Newton (2) Sec
(3) a.m.u. (4) All of these
- Which of the following is not a fundamental unit in SI system ?
(1) ampere (2) candela
(3) kelvin (4) Pascal
- Which one of the following is unitless quantity ?
(1) Velocity gradient
(2) Pressure gradient
(3) Displacement gradient
(4) Force gradient
- What can be the maximum distance of star which can be measured by using parallax method ?
(1) 1 parsec (2) 1 AU
(3) 100 ly (4) Infinite
- Two physical quantities A and B have different dimensions. Which mathematical operation given below is physically possible ?
(1) \sqrt{AB} (2) $A(1+B)$
(3) $A-B$ (4) $A+B$
- Which of the following equations is dimensionally incorrect ?
(E = energy, U = potential energy, P = momentum, m = mass, v = speed)
(1) $E = U + \frac{P^2}{2m}$ (2) $E = mv^2 + \frac{P^2}{m}$
(3) $2E = \frac{U}{2} - \frac{1}{2}mv^2$ (4) $E = \frac{P^2U}{2mv^2}$
- 1 joule of energy is to be converted into new system of units in which length is measured in 10 metre, mass in 10 kg and time in 1 minute. The numerical value of 1 J in the new system is :
(1) 36×10^{-4} (2) 36×10^{-3}
(3) 36×10^{-2} (4) 36×10^{-1}
- The potential energy of a particle varies with distance x from a fixed origin as
 $U = \frac{A\sqrt{x}}{x^2+B}$, where A and B are dimensional constants then dimensional formula for AB is :-
(1) $[ML^{7/2} T^{-2}]$ (2) $[ML^{11/2} T^{-2}]$
(3) $[M^2L^{9/2} T^{-2}]$ (4) $[ML^{13/2} T^{-3}]$
- Which of the following relation cannot be deduced using dimensional analysis ? [the symbols have their usual meanings]
(1) $y = A \sin(\omega t + kx)$ (2) $v = u + at$
(3) $k = \frac{1}{2}mv^2$ (4) All of these
- Of the following quantities which one has dimensions different from the remaining three ?
(1) Energy per unit volume
(2) Force per unit area
(3) Product of voltage and charge per unit volume
(4) Angular momentum per unit mass
- If momentum (P), area (A) and time (T) are taken to be fundamental quantities, then energy has the dimensional formula :
(1) $[P^1A^{-1}T^1]$ (2) $[P^2A^1T^1]$
(3) $[P^1A^{-1/2}T^1]$ (4) $[P^1A^{1/2}T^{-1}]$
- Which of the following specification is most accurate ?
(1) $63.1 \times 10^2 \text{ m}$
(2) $6.31 \times 10^3 \text{ m}$
(3) 6310 m
(4) $0.0631 \times 10^5 \text{ m}$
- If $a = 8 \pm 0.08$ and $b = 6 \pm 0.06$, Let $x = a + b$, $y = a - b$, $z = a \times b$. The correct order of % error in x, y and z
(1) $x = y < z$ (2) $x = y > z$
(3) $x < z < y$ (4) $x > z < y$

16. In an experiment the height of a small object is measured by a Vernier Calliper having least count 0.01 cm is found to be 7.38 cm, then the actual height of object may be :-
 (1) 7.40 cm (2) 7.3942 cm
 (3) 7.3792 cm (4) 7.3882 cm
17. Two resistance are measured in ohm and is given as:-
 $R_1 = 3\Omega \pm 1\%$
 $R_2 = 6\Omega \pm 2\%$
 When they are connected in parallel, the percentage error in equivalent resistance is
 (1) 3% (2) 4.5% (3) 0.67% (4) 1.33%
18. The percentage error in measurement of a physical quantity m given by $m = \pi \tan \theta$ is minimum when
 (1) $\theta = 45^\circ$ (2) $\theta = 90^\circ$
 (3) $\theta = 60^\circ$ (4) $\theta = 30^\circ$
 (Assume that error in θ remain constant)
19. The number of significant digits in 0.001001 is :-
 (1) 6 (2) 4 (3) 7 (4) 2
20. The sides of a rectangle are (10.5 ± 0.2) cm and (5.2 ± 0.1) cm. Then its perimeter with error limits is :-
 (1) (31.4 ± 0.6) cm
 (2) (31.4 ± 0.3) cm
 (3) (51.6 ± 0.6) cm
 (4) (51.6 ± 0.3) cm
21. The Young modulus (Y) of a material is given by $Y = \frac{WL}{\pi r^2 \ell}$. If the percentage error in W, L, r and ℓ are 0.5%, 1%, 3% and 4% respectively then maximum percentage error in Y is -
 (1) 7.5 % (2) 9 % (3) 11.5 % (4) 13 %

ANSWER KEY

Que	1	2	3	4	5	6	7	8	9	10	11	12	13
Ans	3	3	4	4	3	2	1	4	4	2	4	4	4
Que	14	15	16	17	18	19	20	21	22	23	24	25	26
Ans	3	3	3	4	1	2	1	3					

Your Target is to secure Good Rank in Pre-Medical 2019

1. A block of mass 1 kg is placed on a surface for which $\mu_s = 0.2$ & $\mu_k = 0.15$, an external force F is applied on the block as shown in the figure, using $g = 10 \text{ m/sec}^2$, find:

(a) The force of friction on block


(b) Its acceleration

(i) When $F = 1 \text{ N}$


(ii) When $F = 2 \text{ N}$

(iii) When $F = 2.5 \text{ N}$


2. Find value of pulling force F for which block just moves. Given coefficient of friction for surface is μ .




3. Find value of pushing force F for which the body just moves.




4. Find the minimum value of horizontal force (F) required on the block of mass m to keep it at rest on the wall. Given the coefficient of friction between the surfaces is μ .



5. What is the maximum value of force, so that both the blocks will move together?




6.



For the given arrangement, find:


- (i) Force of friction on the block.
 (ii) Acceleration of the block.
 (iii) Minimum Force required to prevent the downward sliding of block.
 (iv) Minimum Force required to just move the block upwards.

7. A block of mass 10 kg is kept over a rough surface and a force $F = 4t$ is applied on it. At what value of t the block will start moving :-




(1) $> 10 \text{ s}$ (2) $< 8 \text{ s}$ (3) $= 9 \text{ s}$ (4) None

8. A block of mass 10 kg is moving on a rough surface as shown in figure. The frictional force acting on block is :-



(1) 20 N (2) 60 N (3) 40 N (4) 80 N

9. A block of mass 10 kg, moving with acceleration 2 m/s^2 on horizontal rough surface is shown in figure



The value of coefficient of kinetic friction is :-

(1) 0.2 (2) 0.4 (3) 0.5 (4) 0.1


10. If acceleration of A is 2 m/s^2 which is smaller than acceleration of B then the value of frictional force applied by B on A is :-

(1) 50 N


(2) 20 N

(3) 10 N

(4) None of these



11. Acceleration of block A varies with time as shown in figure the value of coefficient of kinetic friction between block A and B is




(1) 0.5

(2) 0.6

(3) 0.4


(4) None of these

12. Block of mass 10 kg is moving on inclined plane with constant velocity 10 m/s. The coefficient of kinetic friction between incline plane and block is :-



- (1) 0.57 (2) 0.75
(3) 0.5 (4) None of these


13.



The ratio of acceleration of blocks A placed on smooth incline with block B placed on rough incline is 2 : 1. The coefficient of kinetic friction between block B and incline is :-

- (1) 0.5 (2) 0.75
(3) 0.57 (4) None of these

14.



Blocks shown in figure moves with constant velocity 10 m/s towards right. All surfaces in contact are rough. The friction force applied by B on A is :-

- (1) 0 N
(2) 20 N
(3) 10 N
(4) insufficient data

ANSWERS

1. (i) $f = 1\text{ N}$, $a = 0$

(ii) $f = 2\text{ N}$, $a = 0\text{ m/s}^2$

(iii) $f = 1.5\text{ N}$, $a = 1\text{ m/s}^2$

2. $F = \frac{\mu mg}{\cos\theta + \mu \sin\theta}$

3. $F = \frac{\mu mg}{\cos\theta - \mu \sin\theta}$

4. $F = \frac{mg}{\mu}$

5. $\frac{40}{3}\text{ N}$

6. (i) 12.8 N

(ii) 2.8 m/s²

(iii) 11.2 N

(iv) 36.8 N

7. 1

8. 2

9. 1

10. 3

11. 1

12. 2


13. 1

14. 1

WORK POWER & ENERGY

RACE # 01

1. A block of mass 2 kg is placed on a smooth horizontal surface. Two forces $F_1 = 20$ N and $F_2 = 5$ N start acting on the block in opposite directions as shown. If block gets displaced by 5 m in the direction of net force then work done by F_2 is :-



(1) - 75 J (2) 75 J (3) - 25 J (4) 25 J

2. A man of mass 50 kg is standing in an elevator. If elevator is moving up with an acceleration $\frac{g}{3}$ then work done by normal reaction of elevator floor on man when elevator moves by a distance 12 m is ($g = 10 \text{ m/s}^2$) :-

(1) 2000 J (2) 4000 J (3) 6000 J (4) 8000 J

3. A cubical vessel of height 1 m is full of water. Work done by gravity in taking water-out from vessel will be :-

(1) 5000 J (2) 10,000 J
(3) 5 J (4) 10 J

4. A man pulls a bucket full of water from h metre deep well. If the mass of rope is m and mass of bucket full of water is M , then work done by the man is :-

(1) $\left(\frac{M}{2} + m\right)gh$ (2) $\left(\frac{M+m}{2}\right)gh$

(3) $\left(M + \frac{m}{2}\right)gh$ (4) $(M + m)gh$

5. The potential energy of a particle of mass 1 kg moving along x -axis given

by $U(x) = \left[\frac{x^2}{2} - x\right] \text{ J}$. If total mechanical energy

of particle is 2J then find its maximum speed (in m/s) :-

(1) $\sqrt{5}$ (2) $\sqrt{7}$
(3) $\sqrt{3}$ (4) None

6. The potential energy of an object of mass m moving in xy plane in a conservative field is given by $U = ax + by$, where x and y are position coordinates of the object. Find magnitude of its acceleration :-





(1) $\frac{\sqrt{a^2 + b^2}}{m}$

(2) $\frac{a^2 + b^2}{m}$






(3) $\sqrt{a^2 + b^2}$

(4) None

7. On a particle placed at origin a variable force $F = -ax$ (where a is a positive constant) is applied. If $U(0) = 0$, the graph between potential energy of particle $U(x)$ and x is best represented by :-



8. The variation of potential energy U of a system is shown in figure. The force acting on the system is best represented by :-



9. A particle located in one dimensional potential field has potential energy function

$$U(x) = \frac{a}{x^2} - \frac{b}{x^3}, \text{ where } a \text{ and } b \text{ are positive constants. The position of equilibrium corresponds to } x =$$

(1) $\frac{3a}{2b}$

(2) $\frac{2b}{3a}$

(3) $\frac{2a}{3b}$

(4) $\frac{3b}{2a}$

ANSWER KEY

Ques.	1	2	3	4	5	6	7	8	9
Ans	3	4	1	3	1	1	2	1	4


1. The position (x) of a particle of mass 2 kg moving along x -axis at time t is given by $x = (2t^3)$ metre. Find the work done by force acting on it in time interval $t = 0$ to $t = 2$ s :-

(1) 576 J (2) 584 J
(3) 623 J (4) None

2. The velocity (v) of a particle of mass m moving along x -axis is given by $v = \alpha\sqrt{x}$, where α is a constant. Find work done by force acting on particle during its motion from $x = 0$ to $x = 2m$:-

(1) $m\alpha^2$ (2) $m\alpha$
(3) $\frac{m\alpha}{2}$ (4) None

3. A block of mass 8 kg is released from the top of an inclined smooth surface as shown in figure. If spring constant of spring is 200 N/m and block comes to rest after compressing spring by 1 m then find the distance travelled by block before it comes to rest :-




(1) 2.5 m (2) 3.5 m
(3) 2.0 m (4) None

4. The position (x) of a body moving along x -axis at time (t) is given by $x = 3t^2$, where x is in metre and t is in second. If mass of body is 2 kg, then find the instantaneous power delivered to body by force acting on it at $t = 4$ s :-

(1) 288 W
(2) 280 W
(3) 290 W
(4) None


5. A block of mass m is released on the top of a smooth inclined plane of length x and inclination θ as shown in figure. Horizontal surface is rough. If block comes to rest after moving a distance d on the horizontal surface, then coefficient of friction between block and surface is :-



(1) $\frac{x \sin \theta}{2d}$ (2) $\frac{x \cos \theta}{2d}$
(3) $\frac{x \sin \theta}{d}$ (4) $\frac{x \cos \theta}{d}$

6. A particle of mass 0.1 kg is subjected to a force which varies with distance as shown. If it starts its journey from rest at $x = 0$, then its velocity at $x = 12$ m is :-

(1) 0 m/s
(2) $20\sqrt{2}$ m/s
(3) $20\sqrt{3}$ m/s
(4) 40 m/s



7. An unloaded bus can be stopped by applying brakes on straight road after covering a distance x . Suppose, the passenger add 50% of its weight as the load and the braking force remains unchanged, how far will the bus go after the application of the brakes? (Velocity of bus in both case is same) (Consider negligible friction):-

(1) Zero (2) $1.5x$ (3) $2x$ (4) $2.5x$

8. A body of mass m accelerates uniformly from rest to velocity v_1 in time interval T_1 . The instantaneous power delivered to the body as a function of time t is :-

(1) $\frac{mv_1^2}{T_1^2}t$ (2) $\frac{mv_1}{T_1^2}t$
(3) $\left(\frac{mv_1}{T_1}\right)^2 t$ (4) $\frac{mv_1^2}{T_1}t^2$


9. A car of mass m has an engine which can deliver power P . The minimum time in which car can be accelerated from rest to a speed v is :-

- (1) $\frac{mv^2}{2P}$ (2) Pmv^2
(3) $2Pmv^2$ (4) $\frac{mv^2}{2}P$

10. The rate of doing work by force acting on a particle moving along x -axis depends on position x of particle and is equal to $2x$. The velocity of particle is given by expression :-

- (1) $\left[\frac{3x^2}{m}\right]^{1/3}$ (2) $\left[\frac{3x^2}{2m}\right]^{1/3}$
(3) $\left(\frac{2mx}{9}\right)^{1/2}$ (4) $\left[\frac{mx^2}{3}\right]^{1/2}$

11. An object starts from rest and is acted upon by a variable force F as shown in figure. If F_0 is the initial value of the force, then the position of the object, where it again comes to rest will be :-



- (1) $\frac{2F_0}{\tan \alpha}$ (2) $\frac{F_0}{\sin \alpha}$
(3) $\frac{2F_0}{\cot \alpha}$ (4) $\frac{F_0}{2\cos \alpha}$

ANSWER KEY

Que	1	2	3	4	5	6	7	8	9	10	11
Ans	1	1	1	1	3	4	2	1	1	1	1

CIRCULAR MOTION

on by
 F_0 is
 sition
 be :-

1. A particle of mass m is moving in a circle of radius r under a centripetal force equal to $\frac{-K}{r^2} \hat{r}$,

where K is constant. What is the total energy of the particle ?

- (1) $-K/2r$ (2) $K/4r$
 (3) K/r (4) None

2. If a_r and a_t represents radial and tangential acceleration, the motion of a particle will be circular if :

- (i) $a_r = 0$ and $a_t = 0$ (ii) $a_r = 0$ but $a_t \neq 0$
 (iii) $a_r \neq 0$ but $a_t = 0$ (iv) $a_r \neq 0$ and $a_t \neq 0$
 (1) (i, iii) (2) (ii, iii)
 (3) (iii, iv) (4) (ii, iv)

3. A particle P is moving in a circle of radius ' a ' with uniform speed v . C is the centre of the circle and AB is a diameter. The angular velocity of P about A and C are in the ratio :

- (1) 1 : 1 (2) 1 : 2 (3) 2 : 1 (4) 4 : 1

4. A motor car is travelling at 60 m/s on a circular road of radius 1200 m. It is increasing its speed at the rate of 4 m/s². The acceleration of the car is :

- (1) 3 m/s² (2) 4 m/s²
 (3) 5 m/s² (4) 7 m/s²

5. The speed of a particle moving in a circle slows down at a rate of 3 m/sec². At some instant the magnitude of the total acceleration is 5 m/sec² and the particle speed is 12 m/sec. The radius of circle will be:

- (1) 12 m (2) 24 m (3) 36 m (4) 48 m

5. A point moves along a circle with speed $v = at$. The total acceleration of the point at a time when it has traced 1/8th of the circumference is :

- (1) $\frac{v}{8a}$ (2) $2a\sqrt{4+\pi^2}$

- (3) a (4) $\frac{a}{2}\sqrt{4+\pi^2}$

7. A particle of mass ' m ' is moving along a circle of radius ' r '. At some instant, its speed is ' v ' and it is gaining speed at a uniform rate ' a ', then, at the given instant, acceleration of the particle is :

- (1) along the radius

- (2) inclined to radius at $\theta = \sin^{-1} \frac{1}{\left[1 + \frac{v^4}{a^2 r^2}\right]^{1/2}}$

- (3) inclined to radius at $\theta = \cos^{-1} \frac{ar}{v^2}$

- (4) inclined to radius at $\theta = \tan^{-1} \frac{v^2}{ar}$

8. Particle A and B are moving in coplanar circular paths centred at O . They are rotating in the same sense. Time periods of rotation of A and B around O are T_A and T_B , respectively, with $T_B > T_A$. Time required for B to make one rotation around O relative to A is :

- (1) $T_B - T_A$

- (2) $T_B + T_A$

- (3) $\frac{T_B T_A}{T_B + T_A}$

- (4) $\frac{T_B T_A}{T_B - T_A}$



9. For a particle in uniform circular motion, the acceleration \vec{a} at a point $P(R, \theta)$ on the circle of radius R is (Here θ is measured from the x -axis) :

- (1) $\frac{v^2}{R} \hat{i} + \frac{v^2}{R} \hat{j}$

- (2) $-\frac{v^2}{R} \cos \theta \hat{i} + \frac{v^2}{R} \sin \theta \hat{j}$

- (3) $-\frac{v^2}{R} \sin \theta \hat{i} + \frac{v^2}{R} \cos \theta \hat{j}$

- (4) $-\frac{v^2}{R} \cos \theta \hat{i} - \frac{v^2}{R} \sin \theta \hat{j}$

Together, we will make a difference.



Sitting: Govind Maheshwari (Director), Rajesh Maheshwari (Director)
Standing: Naveen Maheshwari (Director), Brajesh Maheshwari (Director & H.O.D. Physics)

ALLEN Career Institute

Corporate Office : "SANKALP", CP-6, Indra Vihar, Kota (Rajasthan) INDIA 324005 Ph: +91-744-5156100 e-mail: info@allen.ac.in

AHMEDABAD H.O.

Off. S.G. Highway, Behind Patel Avenue
Near Purshottam Bungalow
Bodakdev, Ahmedabad (Gujarat)
Tel: 079-40306001

BENGALURU H.O.

No. 36, 15th Cross, 3rd Block, New South
End Circle, Opp. Central Library,
Jayanagar, Bengaluru
Tel: 080-46704000

CHANDIGARH H.O.

SCO-354-355
Ground Floor,
Sector 34-A Chandigarh (UT) 160022
Tel: 0172-5257100

INDORE H.O.

Agrawal Palace (Plote No. 2)
Opp. Bhoaskar Hospital
Behind High Court, Nath Mandir Road
South Tukoganj, INDORE
Tel: 0731-4728500

JAIPUR H.O.

11, Samachar Jagat Building,
Opposite Vidyasharam School,
JLN Marg, Jaipur
Tel: 0141-5195100

MUMBAI H.O.

5th Floor, Baba house, Western Express
Highway Metro Station, Andheri-Kurla Road,
andheri East, Mumbai
Tel: 022-40040602