



# CHEMISTRY

Target : JEE (Main)

QUALITATIVE ANALYSIS

WWW.GRADESETTER.COM



# QUALITATIVE ANALYSIS

## CONTENTS

Topic	Page No.
Theory	01 – 21
Exercise - 1	22 – 26
Exercise - 2	27 – 28
Exercise - 3	28
JEE (Main) /AIEEE Questions	
Answer Key	29

### JEE (MAIN) SYLLABUS

**Chemical Principle involved in the qualitative salt analysis :**

Cations -  $\text{Pb}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Al}^{3+}$ ,  $\text{Fe}^{3+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{NH}_4^+$ .

Anions -  $\text{CO}_3^{2-}$ ,  $\text{S}^{2-}$ ,  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$ ,  $\text{NO}_2^-$ ,  $\text{Cl}^-$ ,  $\text{Br}^-$ ,  $\text{I}^-$ . (Insoluble salts excluded).

### JEE(ADVANCED) SYLLABUS

**Principles of Qualitative Analysis :** Groups I to V (only  $\text{Ag}^+$ ,  $\text{Hg}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Bi}^{3+}$ ,  $\text{Fe}^{3+}$ ,  $\text{Cr}^{3+}$ ,  $\text{Al}^{3+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Mn}^{2+}$  and  $\text{Mg}^{2+}$ ); Nitrate, halides (excluding fluoride), sulphate and sulphide.

© Copyright reserved.

All rights reserved. Any photocopying, publishing or reproduction of full or any part of this study material is strictly prohibited. This material belongs to an enrolled student of RESONANCE only any sale/resale of this material is punishable under law, subject to Kota Jurisdiction only.



# Qualitative Analysis

## Introduction :

Qualitative analysis involves the detection of cation(s) and anion(s) of a salt or a mixture of salts. The systematic procedure for qualitative analysis of an inorganic salt involves the following steps :

### (a) Preliminary tests

- Physical appearance (colour and smell).
- Flame test.
- Dilute sulphuric acid test.
- Concentrated sulphuric acid test.
- Dry heating test.
- Borax bead test.
- Potassium permanganate test.
- Tests for sulphate, phosphate and borate.

### (b) Wet tests for acid radicals.

### (c) Wet tests (group analysis) for basic radicals.

#### 1. Physical Examination of the Mixture :

The physical examination of the unknown mixture involves the study of colour, smell and density.


#### 2. Dry Heating Test :

This test is performed by heating a small amount of mixture in a dry test tube. Quite valuable information can be generated by carefully performing and noting the observations here. On heating some salts undergo decomposition thus evolving the gases or may undergo characteristic changes in the colour of residue.


#### 3. Flame test :

The chlorides of the metals are more volatile as compared to other salts and these are prepared in situ by mixing the compounds with a little concentrated hydrochloric acid. On heating in a non-luminous Bunsen flame they are volatilized and impart a characteristic colour to the flame as these absorb energy from the flame and transmit the same as light as characteristic colour .

Colour of Flame	Inference
Crimson Red / Carmine Red	Lithium
Golden yellow	Sodium
Violet/Lilac	Potassium
Brick red	Calcium
Crimson	Strontium
Apple Green/Yellowish Green	Barium
Green with a Blue centre/Greenish Blue	Copper



(A) Dipping the platinum wire in the paste of salt and HCl.




(B) Introducing the wire in the flame






4. **Borax Bead test :**

On heating borax forms a colourless glassy bead of  $\text{NaBO}_2$  and  $\text{B}_2\text{O}_3$ .





On heating with a coloured salt, the glassy bead forms a coloured metaborate in oxidising flame.

For example, in oxidising flame copper salts give blue bead.



However, in reducing flame the colours may be different due to different reactions.




Metal	Colour in oxidising flame		Colour in reducing flame	
	When Hot	When Cold	When Hot	When Cold
Copper	Green	Blue	Colourless	Brown red
Iron	Brown yellow	Pale yellow/Yellow	Bottle green	Bottle green
Chromium	Yellow	Green	Green	Green
Cobalt	Blue	Blue	Blue	Blue
Manganese	Violet/Amethyst	Red/Amethyst	Grey/Colourless	Grey/Colourless
Nickel	Violet	Brown/Reddish brown	Grey	Grey

**Analysis of ANIONS (Acidic Radicals) :**


**(a) DILUTE SULPHURIC ACID/DILUTE HYDROCHLORIC ACID GROUP :**

**1. CARBONATE ION ( $\text{CO}_3^{2-}$ ):**


- Dilute  $\text{H}_2\text{SO}_4$  test : A colourless odourless gas is evolved with brisk effervescence.




- Lime water/Baryta water ( $\text{Ba}(\text{OH})_2$ ) test : The liberated gas can be identified by its property of rendering lime water (or baryta water) turbid.



On prolonged passage of  $\text{CO}_2$  the milkiness disappears.




- ☞ Soluble bicarbonates give white precipitate with  $\text{MgSO}_4$  (aq) /  $\text{MgCl}_2$  (aq) only on heating.




**2. SULPHITE ION ( $\text{SO}_3^{2-}$ ):**


- Dilute  $\text{H}_2\text{SO}_4$  test : Decomposition of salt is more rapidly on warming, with the evolution of sulphur dioxide.



- Acidified potassium dichromate test : The filter paper dipped in acidified  $\text{K}_2\text{Cr}_2\text{O}_7$  turns green.





- Barium chloride/Strontium chloride solution : White precipitate of barium (or strontium) sulphite is obtained.



- ☞ White precipitate is soluble in dilute HCl forming  $\text{SO}_2$ .



- Lime water test : A white precipitate is formed. The precipitate dissolves on prolonged passage of the gas, due to the formation of hydrogen sulphite ions.





Qualitative Analysis

**3. SULPHIDE ION (S<sup>2-</sup>):**

- **Dilute H<sub>2</sub>SO<sub>4</sub> test:** Pungent smelling gas like that of rotten egg is obtained.  

$$S^{2-} + 2H^+ \longrightarrow H_2S \uparrow$$
- **Lead acetate test:** Filter paper moistened with lead acetate solution turns black.  

$$(CH_3COO)_2Pb + H_2S \longrightarrow PbS \downarrow (\text{black}) + 2CH_3COOH.$$
- **Sodium nitroprusside test:** Purple coloration is obtained.  

$$S^{2-} + [Fe(CN)_5(NO)]^{2-} \longrightarrow [Fe(CN)_5NOS]^{4-} (\text{violet}).$$
- **Cadmium carbonate suspension/ Cadmium acetate solution:** Yellow precipitate is formed.  

$$Na_2S + CdCO_3 \longrightarrow CdS \downarrow + Na_2CO_3$$

**4. NITRITE ION (NO<sub>2</sub><sup>-</sup>):**

- **Dilute H<sub>2</sub>SO<sub>4</sub> test:** Solid nitrite in cold produces a transient pale blue liquid (due to the presence of free nitrous acid, HNO<sub>2</sub> or its anhydride, N<sub>2</sub>O<sub>3</sub>) first and then evolution of pungent smelling reddish brown vapours of NO<sub>2</sub> takes place.  

$$NO_2^- + H^+ \longrightarrow HNO_2; (2HNO_2 \longrightarrow H_2O + N_2O_3);$$

$$3HNO_2 \longrightarrow HNO_3 + 2NO + H_2O; 2NO + O_2 \longrightarrow 2NO_2 \uparrow$$
- **Starch iodide test:** The addition of a nitrite solution to a solution of potassium iodide, followed by acidification with acetic acid or with dilute sulphuric acid, results in the liberation of iodine, which may be identified by the blue colour produced with starch paste. A similar result is obtained by dipping potassium iodide-starch paper moistened with a little dilute acid into the solution.  

$$2NO_2^- + 3I^- + 4CH_3COOH \longrightarrow I_3^- + 2NO \uparrow + 4CH_3COO^- + 2H_2O$$

$$\text{Starch} + I_3^- \longrightarrow \text{Blue (starch iodine adsorption complex)}$$
- **Ferrous sulphate test (Brown ring test):** When the nitrite solution is added carefully to a concentrated solution of iron(II) sulphate acidified with dilute acetic acid or dilute sulphuric acid, a brown ring appears due to the formation of [Fe(H<sub>2</sub>O)<sub>5</sub>NO]SO<sub>4</sub> at the junction of the two liquids. If the addition has not been made slowly and cautiously, a brown colouration results.  

$$NO_2^- + CH_3COOH \longrightarrow HNO_2 + CH_3COO^-$$

$$3HNO_2 \longrightarrow H_2O + HNO_3 + 2NO \uparrow$$

$$Fe^{2+} + SO_4^{2-} + NO \uparrow \longrightarrow [Fe, NO]SO_4$$
- **Thiourea test:** When a dilute acetic acid solution of a nitrite is treated with a little solid thiourea, nitrogen is evolved and thiocyanic acid is produced. The latter may be identified by the red colour produced with dilute HCl and FeCl<sub>3</sub> solution.  

$$NaNO_2 + CH_3COOH \longrightarrow HNO_2 + CH_3COONa$$

$$HNO_2 + H_2NCSNH_2(s) \text{ (thiourea)} \longrightarrow N_2 + HSCN + 2H_2O$$

$$FeCl_3 + 3HSCN \xrightarrow{\text{dil HCl}} Fe(SCN)_3 \text{ (blood red colouration)} + 3HCl$$

**5. ACETATE ION (CH<sub>3</sub>COO<sup>-</sup>)**

- With dilute H<sub>2</sub>SO<sub>4</sub> a vinegar like smell is obtained.  

$$(CH_3COO)_2Ca + H_2SO_4 \longrightarrow 2CH_3COOH + CaSO_4$$
  - **Neutral ferric chloride test:** A deep red/ blood red colouration (no precipitate) indicates the presence of acetate.  

$$6CH_3COO^- + 3Fe^{3+} + 2H_2O \longrightarrow [Fe_3(OH)_2(CH_3COO)_6]^{+} + 2H^+$$
- When solution is diluted with water and boiled, brownish red precipitate of basic iron (III) acetate is obtained.
- $$[Fe_3(OH)_2(CH_3COO)_6]^{+} + 4H_2O \xrightarrow{\text{Boil}} 3Fe(OH)_2CH_3COO^- + 3CH_3COOH + H^+$$

Solved Examples

**Ex.1** An aqueous solution of salt containing an anion X<sup>n-</sup> gives the following reactions:  
 (i) It gives the purple or violet colouration with sodium nitroprusside solution.  
 (ii) It liberates a colourless unpleasant smelling gas with dilute H<sub>2</sub>SO<sub>4</sub> which turns lead acetate paper black.  
 Identify the anion (X<sup>n-</sup>) and write the chemical reactions involved.




Qualitative Analysis

- Sol.**  $X^{n-}$  is  $S^{2-}$  because
- (i)  $[Fe(CN)_6NO]^{2-} + S^{2-} \longrightarrow [Fe(CN)_6NOS]^{4-}$  (purple or violet colouration)
- (ii)  $S^{2-} + H_2SO_4 \longrightarrow H_2S \uparrow$  (colourless unpleasant smelling) +  $SO_4^{2-}$
- $H_2S + Pb(CH_3COO)_2 \longrightarrow PbS \downarrow$  (black) +  $2CH_3COOH$
- Ex.2** Sulphite on treatment with dil.  $H_2SO_4$  liberates a gas which :
- (1) turns lead acetate paper black (2) burns with blue flame  
 (3) smells like vinegar (4) turns acidified  $K_2Cr_2O_7$  solution green
- Sol.**  $SO_3^{2-} + H_2SO_4 \longrightarrow SO_2 + SO_4^{2-} + H_2O$   
 $SO_2$  turns acidified  $K_2Cr_2O_7$  solution green.  
 $K_2Cr_2O_7 + H_2SO_4 + 3SO_2 \longrightarrow Cr_2(SO_4)_3$  (Green) +  $K_2SO_4 + H_2O$   
 Therefore, (4) option is correct.


(b) CONC.  $H_2SO_4$  GROUP :

1. CHLORIDE ION ( $Cl^-$ ) :


- **Concentrated  $H_2SO_4$  test :** Colourless pungent smelling gas is evolved which gives fumes of  $NH_4Cl$  when a glass rod dipped in dil.  $HCl$  is brought in contact with evolving gas.




- $NH_4OH + HCl \longrightarrow NH_4Cl \uparrow$  (white fumes) +  $H_2O$ .




- **Silver nitrate test :**





- ☞ White precipitate is soluble in aqueous ammonia and precipitate reappears with  $HNO_3$ .



- **Chromyl Chloride test :**




When deep red vapours are passed into sodium hydroxide solution, a yellow solution of sodium chromate is formed, which when treated with lead acetate gives yellow precipitate of lead chromate.






- ☞ **Heavy metal chlorides** such as  $Hg_2Cl_2$ ,  $HgCl_2$ ,  $SnCl_2$ ,  $AgCl$ ,  $PbCl_2$  and  $SbCl_3$  do not respond to this test as they are partially dissociated. This test is given generally by **ionic chlorides**.

- ☞ Test should be carried out in a dry test tube otherwise chromic acid will be formed.




2. BROMIDE ION ( $Br^-$ ) :

- **Concentrated  $H_2SO_4$  test :** First a reddish-brown solution is formed, then reddish-brown bromine vapour accompanies the hydrogen bromide (fuming in moist air) is evolved.




- **Silver Nitrate test :** Pale yellow precipitate is formed



- ☞ Yellow precipitate is partially soluble in dilute aqueous ammonia but readily dissolves in concentrated ammonia solution.



- **Chlorine water test (organic layer test) :** When to a sodium carbonate extract of metal bromide containing  $CCl_4$ ,  $CHCl_3$  or  $CS_2$ , chlorine water is added and the content is shaken and then allow to settle down reddish brown colour is obtained in organic layer.






**Qualitative Analysis**


$Br_2 + CHCl_3 / CCl_4 \longrightarrow Br_2$  dissolve to give reddish brown colour in organic layer.  
 With excess of chlorine water, the bromine is converted into yellow bromine monochloride and a pale yellow solution results.  
 $Br_2 \uparrow + Cl_2 \uparrow \longrightarrow 2BrCl$

**3. IODIDE ION ( $I^-$ ):**


**Concentrated  $H_2SO_4$  test:** Pungent smelling violet vapours are evolved.



Evolution of dark violet fumes intensifies on adding a pinch of  $MnO_2$ .



**Starch paper test:** Iodides are readily oxidised in acid solution to free iodine; the free iodine may than be identified by deep blue colouration produced with starch solution.





**Silver nitrate test:** Bright yellow precipitate is formed.




Bright yellow precipitate is insoluble in dilute aqueous ammonia but is partially soluble in concentrated ammonia solution.

**Chlorine water test (organic layer test):** When chlorine water is added to a solution of iodide, free iodine is liberated which colours the solution brown and on shaking with  $CS_2$ ,  $CHCl_3$  or  $CCl_4$ , it dissolves in organic layer forming a violet solution, which settles below the aqueous layer.




If excess of chlorine water is added,  $I_2$  is oxidised to iodic acid (colourless).





**4. NITRATE ION ( $NO_3^-$ ):**





**Concentrated  $H_2SO_4$  test:** Pungent smelling reddish brown vapours are evolved.



Addition of bright copper turnings or paper pellets intensifies the evolution of reddish brown gas.



**Brown ring test:** When a freshly prepared saturated solution of iron (II) sulphate is added to nitrate solution and then concentrated  $H_2SO_4$  is added slowly from the side of the test tube, a brown ring is obtained at the junction of two layers.



On shaking and warming the mixture,  $NO$  escapes and a yellow solution of iron(III) ions is obtained.

**(B) GROUP 'B' RADICALS:**


Group of anions which do not give any gas with dilute as well as concentrated  $H_2SO_4$  in cold but give precipitate with certain reagents:

These acid radicals are identified in inorganic salts by their individual tests as given below

**1. SULPHATE ION ( $SO_4^{2-}$ ):**

**Barium chloride test:**

W.E. or S.E. + Barium chloride (aq)  $\longrightarrow$  White precipitate





**Qualitative Analysis**

- ☞ White precipitate is insoluble in warm dil.  $\text{HNO}_3$  as well as  $\text{HCl}$  but moderately soluble in boiling concentrated hydrochloric acid.
- **Lead acetate test :**  
 $\text{W.E. or S.E.} + \text{Lead acetate} \longrightarrow \text{white precipitate}$   
 $\text{Na}_2\text{SO}_4 + (\text{CH}_3\text{COO})_2\text{Pb} \longrightarrow \text{PbSO}_4 \downarrow (\text{White}) + 2\text{CH}_3\text{COONa}$
- ☞ White precipitate soluble in excess of hot ammonium acetate.  
 $\text{PbSO}_4 + 2\text{CH}_3\text{COONH}_4 \longrightarrow (\text{CH}_3\text{COO})_2\text{Pb (soluble)} + (\text{NH}_4)_2\text{SO}_4$
- **Match stick test :**  
 (a)  $\text{W.E. or S.E.} + \text{Barium chloride} \longrightarrow \text{white precipitate}$   
 $\text{Na}_2\text{SO}_4 + \text{BaCl}_2 \longrightarrow 2\text{NaCl} + \text{BaSO}_4 \downarrow (\text{white})$   
 (b) White precipitate +  $\text{Na}_2\text{CO}_3(\text{s})$  mix and apply the paste on the end of the carbonized match stick or a wooden splinter. Put it in the reducing flame.  
 $\text{BaSO}_4 (\text{s}) + \text{Na}_2\text{CO}_3(\text{s}) \longrightarrow \text{Na}_2\text{SO}_4 + \text{BaCO}_3 \downarrow (\text{white})$   
 $\text{Na}_2\text{SO}_4 + 4\text{C} \longrightarrow \text{Na}_2\text{S} + 4\text{CO}$   
 (c) Now dip the match stick in sodium nitroprusside solution, purple colour near the fused mass is developed.  
 $\text{Na}_2\text{S} + \text{Na}_2[\text{Fe}(\text{CN})_5\text{NO}] \longrightarrow \text{Na}_4[\text{Fe}(\text{CN})_5\text{NOS}] (\text{purple})$

**2. PHOSPHATE ION ( $\text{PO}_4^{3-}$ ) :**

- **Ammonium molybdate test :**  
 $\text{Na}_2\text{HPO}_4 (\text{aq}) + 12(\text{NH}_4)_2\text{MoO}_4 + 23\text{HNO}_3 \longrightarrow (\text{NH}_4)_3\text{PMo}_{12}\text{O}_{40} \downarrow (\text{canary yellow}) + 2\text{NaNO}_3 + 21\text{NH}_4\text{NO}_3 + 12\text{H}_2\text{O}$
- ☞ Some times ammonium phosphomolybdate is also represented by the formula  $(\text{NH}_4)_3\text{PO}_4 \cdot 12\text{MoO}_3$
- **Magnesium nitrate or magnesia mixture test :**  $\text{W.E. or S.E.} + \text{Magnesium nitrate reagent}$  (3-4 mL) and allows to stand for 4-5 minutes, white crystalline precipitate is formed.  
 $\text{Na}_2\text{HPO}_4 (\text{aq}) + \text{Mg}(\text{NO}_3)_2 (\text{aq}) + \text{NH}_4\text{OH}(\text{aq}) \longrightarrow \text{Mg}(\text{NH}_4)\text{PO}_4 \downarrow (\text{white}) + 2\text{NaNO}_3 + \text{H}_2\text{O}$   
 Magnesia mixture is a solution containing  $\text{MgCl}_2$ ,  $\text{NH}_4\text{Cl}$  and a little aqueous  $\text{NH}_3$ .
- ☞  $\text{PO}_4^{3-}$  also gives  $\text{BaCl}_2$  test due to the formation of white precipitate of  $\text{Ba}_3(\text{PO}_4)_2$ . So phosphate test should be carried out first and then conclude if  $\text{PO}_4^{3-}$  is present or absent before proceeding with the test for  $\text{SO}_4^{2-}$ .
- **Silver nitrate solution :** Yellow precipitate is formed which is soluble in dilute ammonia and in dilute nitric acid.  
 $\text{PO}_4^{3-} + 3\text{Ag}^+ \longrightarrow \text{Ag}_3\text{PO}_4 \downarrow$   
 $\text{Ag}_3\text{PO}_4 \downarrow + 6\text{NH}_3 \longrightarrow 3[\text{Ag}(\text{NH}_3)_2]^+ + \text{PO}_4^{3-}$  ;  $\text{Ag}_3\text{PO}_4 \downarrow + 2\text{H}^+ \longrightarrow \text{H}_2\text{PO}_4^- + 3\text{Ag}^+$

**3. BORATE ION ( $\text{BO}_3^{3-}$ ) :**

- ☞ Salt (0.2 g) + conc.  $\text{H}_2\text{SO}_4$  (1 mL) + Ethyl alcohol (4-5 mL) mix in a test tube and then heat. Ignite the evolved vapours with the help of Bunsen flame, green edged flame is obtained.  
 $\text{Na}_3\text{BO}_3 + 3\text{H}_2\text{SO}_4 \longrightarrow 3\text{Na}_2\text{SO}_4 + 2\text{H}_3\text{BO}_3$   
 $3\text{C}_2\text{H}_5\text{OH} + \text{H}_3\text{BO}_3 \longrightarrow (\text{C}_2\text{H}_5)_3\text{BO}_3 + 3\text{H}_2\text{O}$

**Solved Examples**

**Ex.3** A compound (A) of S, Cl and O has vapour density of 67.5 (approx.). It reacts with water to form two acids and reacts with  $\text{KOH}$  to form two salts (B) and (C) while (B) gives white precipitate with  $\text{AgNO}_3$  solution and (C) gives white precipitate with  $\text{BaCl}_2$  solution. Identify (A), (B) & (C).

**Sol.** As mixture give white precipitate with  $\text{BaCl}_2$  and  $\text{AgNO}_3$ , it should contain  $\text{SO}_4^{2-}$  and  $\text{Cl}^-$  ions. As  $\text{SO}_2\text{Cl}_2$  when dissolved in water gives, a mixture of  $\text{H}_2\text{SO}_4$  &  $\text{HCl}$  which then react with  $\text{KOH}$  to form  $\text{KCl}$  and  $\text{K}_2\text{SO}_4$ . Therefore, (A) is  $\text{SO}_2\text{Cl}_2$  and (B) & (C) are  $\text{K}_2\text{SO}_4$  and  $\text{KCl}$  respectively.

Vapour density of  $\text{SO}_2\text{Cl}_2$  = molecular weight / 2.

Vapour density of  $\text{SO}_2\text{Cl}_2$  =  $135 / 2 = 67.5$ .

**Qualitative Analysis**

**Ex.4** Bromine var (1) brown  
**Sol.**  $2\text{I}^- + \text{Br}_2 \longrightarrow$   
 Therefore,

**Ex.5**  $\text{Na}_2\text{S}_2\text{O}_3$   
 (1)  $\text{Na}_2\text{S}_2\text{O}_3$   
**Sol.**  $2\text{Na}_2\text{S}_2\text{O}_3$   
 Therefore

**Analysis of**

Group

Zn



Qualitative Analysis

- Ex.4** Bromine vapours turn moist starch iodide paper :  
 (1) brown (2) red (3) blue (4) colourless  
**Sol.**  $2I^- + Br_2 \rightarrow I_2 + 2Br^-$ ;  $I_2 + \text{starch} \rightarrow \text{blue starch iodine adsorption complex}$ .  
 Therefore, (3) option is correct.
- Ex.5**  $Na_2S_2O_3 + I_2 \rightarrow NaI + \dots [X]$ , [X] is :  
 (1)  $Na_2S_4O_6$  (2)  $Na_2SO_4$  (3)  $Na_2S$  (4)  $Na_2SO_3$   
**Sol.**  $2Na_2S_2O_3 + I_2 \rightarrow 2NaI + Na_2S_4O_6$ .  
 Therefore, (1) option is correct.

**Analysis of CATIONS (Basic Radicals) :**

Group	Group reagent	Basic radical	Composition and colour of precipitate
Zero	NaOH or Ca(OH) <sub>2</sub> , heat if required	NH <sub>4</sub> <sup>+</sup>	Ammonia gas is evolved.
1.	Dil HCl	Ag <sup>+</sup> Hg <sub>2</sub> <sup>2+</sup> Pb <sup>2+</sup>	AgCl ; White Hg <sub>2</sub> Cl <sub>2</sub> ; White PbCl <sub>2</sub> ; White
2.(A)	H <sub>2</sub> S in presence of dil HCl (Insoluble in YAS)	Hg <sup>2+</sup> Pb <sup>2+</sup> Bi <sup>3+</sup> Cu <sup>2+</sup> Cd <sup>2+</sup>	HgS ; Black PbS ; Black Bi <sub>2</sub> S <sub>3</sub> ; Black CuS ; Black CdS ; Yellow
2.(B)	H <sub>2</sub> S in presence of dil HCl (Soluble in YAS)	As <sup>3+</sup> Sb <sup>3+</sup> Sn <sup>2+</sup> Sn <sup>4+</sup>	As <sub>2</sub> S <sub>3</sub> ; Yellow Sb <sub>2</sub> S <sub>3</sub> ; Orange SnS ; Brown SnS <sub>2</sub> ; Yellow
3.	NH <sub>4</sub> OH in presence of NH <sub>4</sub> Cl	Fe <sup>3+</sup> Cr <sup>3+</sup> Al <sup>3+</sup>	Fe(OH) <sub>3</sub> ; Reddish brown Cr(OH) <sub>3</sub> ; Green Al(OH) <sub>3</sub> ; Gelatinous white
4.	H <sub>2</sub> S in presence of NH <sub>4</sub> OH and NH <sub>4</sub> Cl	Zn <sup>2+</sup> Mn <sup>2+</sup> Co <sup>2+</sup> Ni <sup>2+</sup>	ZnS ; White MnS ; Buff (or Pink) CoS ; Black NiS ; Black
5.	(NH <sub>4</sub> ) <sub>2</sub> CO <sub>3</sub> in presence of NH <sub>4</sub> OH	Ba <sup>2+</sup> Sr <sup>2+</sup> Ca <sup>2+</sup>	BaCO <sub>3</sub> ; White SrCO <sub>3</sub> ; White CaCO <sub>3</sub> ; White
6.	Na <sub>2</sub> HPO <sub>4</sub> in presence of NH <sub>4</sub> OH [YAS = Yellow ammonium sulphide. (NH <sub>4</sub> ) <sub>2</sub> S <sub>x</sub> ].	Mg <sup>2+</sup>	Mg(NH <sub>4</sub> )PO <sub>4</sub> ; White


**ZERO GROUP :**

**1. AMMONIUM ION (NH<sub>4</sub><sup>+</sup>) :**

**Sodium hydroxide solution :** Ammonia gas is evolved on warming the solution containing ammonium salt and sodium hydroxide.



- ☞ The gas can be identified by the following characteristics / reactions.
- Its characteristics smell.
- The evolution of the white fumes of ammonium chloride when a glass rod dipped in dilute HCl is held in the vapour.






Qualitative Analysis

- Its ability to turn filter paper moistened with  $Hg_2(NO_3)_2$  solution black.  
 $2HgNO_3 + 2NH_3 \longrightarrow \underset{\text{black}}{Hg(NH_2)NO_3 + Hg} + NH_4NO_3$
- Its ability to turns filter paper moistened with  $CuSO_4$  solution deep blue.  
 $CuSO_4 + 4NH_3 \longrightarrow [Cu(NH_3)_4]SO_4$
- **Nessler's reagent (Alkaline solution of potassium tetraiodomercurate(II)) :**  
 Brown precipitate or brown or yellow colouration is obtained according to the amount of ammonia or ammonium ions present. The precipitate is a basic mercury (II) amido-iodide.  
 $NH_4^+ + 2[HgI_4]^{2-} + 4OH^- \longrightarrow HgO \cdot Hg(NH_2)I \downarrow + 7I^- + 3H_2O$

**1<sup>st</sup> GROUP ( $Pb^{2+}$ ,  $Hg_2^{2+}$ ,  $Ag^+$ ) :**



Qualitative An

**1. LEAD ION (**

- Dilut
- White KCl.
- Sc
- F
- Ye
- 
- f


**2. ME**




Qualitative Analysis

**1. LEAD ION (Pb<sup>2+</sup>) :**


• **Dilute HCl solution :** White precipitate is formed in cold solution.




White precipitate is soluble in hot water. White precipitate is also soluble in concentrated HCl or concentrated KCl.



• **Sodium hydroxide solution :** White precipitate is formed which is soluble in excess of the reagent.



• **Potassium iodide solution :** A yellow precipitate is formed.




Yellow precipitate reappears on dilution with water.

• **Potassium chromate solution (in neutral, acetic acid or ammonia solution) :** A yellow precipitate is formed.




Yellow precipitate is soluble in sodium hydroxide and HNO<sub>3</sub> (nitric acid).

• **Ammonia solution :** With ammonia solution, Pb<sup>2+</sup> gives a white precipitate of lead hydroxide.




**2. MERCURY (I) ION (Hg<sub>2</sub><sup>2+</sup>) :**


• **Dilute HCl solution :** White precipitate is formed in cold solution.




• **Ammonia solution :** A mixture of mercury metal (black precipitate) and basic mercury (II) amido chloride (white precipitate) is formed.




• **Dissolution of white precipitate (Hg<sub>2</sub>Cl<sub>2</sub>) in aquaregia :**




(a) **Stannous chloride test :** White precipitate is formed which finally turns to black.




(b) **Potassium iodide test :** Scarlet/red precipitate is formed which is soluble in excess of the reagent.



(c) **Copper chips test :** Shining grey deposition of mercury on copper chips is formed.



• **Reaction with H<sub>2</sub>S gas :**



**3. SILVER ION (Ag<sup>+</sup>) :**

• **Dilute hydrochloric acid/soluble chlorides :** White precipitate is formed.




The precipitate obtained after filtration is soluble in concentrated HCl.




Dilute ammonia solution dissolves the precipitate forming a soluble complex.




Dilute nitric acid or hydrochloric acid neutralizes the excess ammonia and the precipitate reappears because the equilibrium is shifted backwards.



• **Potassium iodide solution :** A bright yellow precipitate is formed which is insoluble in dilute ammonia but partially soluble in concentrated ammonia.



The yellow precipitate is soluble in KCN and in Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>.





Qualitative Analysis


● Potassium chromate solution: Red precipitate is formed which is soluble in dilute HNO<sub>3</sub> and in ammonia solution.  
 $2Ag^+ + CrO_4^{2-} \rightarrow Ag_2CrO_4 \downarrow$

**II<sup>nd</sup> Group (Hg<sup>2+</sup>, Pb<sup>2+</sup>, Bi<sup>3+</sup>, Cu<sup>2+</sup>, Cd<sup>2+</sup>, As<sup>3+</sup>, Sb<sup>3+</sup>, Sn<sup>2+</sup>)**

On the basis of the solubility of the precipitates of the sulphides of the II group cations in yellow ammonium sulphide, they have been classified into two subgroups as given below:

- IIA :** HgS, PbS, CuS, Bi<sub>2</sub>S<sub>3</sub>, all black but CdS is yellow. All insoluble in yellow ammonium sulphide.  
**IIB :** SnS<sub>2</sub>, As<sub>2</sub>S<sub>3</sub> are yellow, Sb<sub>2</sub>S<sub>3</sub> is orange & SnS is dark brown All soluble in yellow ammonium sulphide.

**IIA Group (Hg<sup>2+</sup>, Pb<sup>2+</sup>, Bi<sup>3+</sup>, Cu<sup>2+</sup>, Cd<sup>2+</sup>)**




2.




Qualitative Analysis

1. MERCURY (II) ION ( $Hg^{2+}$ ) :


● **Precipitation with  $H_2S$  in acidic medium** : Black precipitate is formed. Precipitate insoluble in water, hot dilute  $HNO_3$ , alkali hydroxides, or colourless ammonium sulphide.




☞ Aqua regia dissolves the precipitate.



● **Potassium iodide solution** : On slow addition red precipitate is formed.




Precipitate dissolves in excess of KI forming colourless soluble complex.



● **Ammonia solution** : White precipitate of mixed composition (Mercury (II) oxide + Mercury (II) amido nitrate) is formed with metal nitrate.




● **Cobalt (II) thiocyanate test** : When reagent is added to an aqueous solution of  $Hg^{2+}$  ions and the walls of the test tube is stirred with a glass rod, deep-blue crystalline precipitate is formed.



☞ In place of Cobalt (II) thiocyanate,  $Co(CH_3COO)_2$  and  $NH_4SCN$  can be added to the aqueous solution of  $Hg^{2+}$  ions.


2. COPPER ION ( $Cu^{2+}$ ) :

● **Precipitation with  $H_2S$  in acidic medium** : Black precipitate is formed.




Precipitate is insoluble in boiling dilute (M)  $H_2SO_4$  (distinction from cadmium), in  $NaOH$ ,  $Na_2S$  and  $(NH_4)_2S$ .

Precipitate dissolves in hot concentrated  $HNO_3$




When boiled for longer S is oxidised to  $H_2SO_4$  and a clear solution of  $Cu(NO_3)_2$  is obtained.


● **Ammonia solution** : When added sparingly a blue precipitate of basic salt (basic copper sulphate) is formed with  $CuSO_4$ .




It is soluble in excess of reagent forming a deep blue colouration.



● **Sodium hydroxide in cold solution** : A blue precipitate is formed.



● **Potassium iodide** : It gives a white precipitate of  $Cu(I)$  iodide but the solution is intensely brown because of the formation of tri-iodide ions (or iodine).



The solution becomes colourless and a white precipitate is visible when excess of sodium thiosulphate solution is added.



☞ These reactions are used in quantitative analysis for the iodometric determination of copper.


● **Potassium ferrocyanide (Potassium hexacyanidoferrate (II)) solution** :  $Cu^{2+}$  ions gives brown/chocolate brown precipitate.






Qualitative Analysis


(vi) **Potassium cyanide** : When added sparingly forms first a yellow precipitate.



Precipitate quickly decomposes into CuCN and cyanogen.






Excess reagent dissolves the precipitate forming a colourless soluble complex.




Complex is so stable that H<sub>2</sub>S cannot precipitate Cu (I) sulphide (distinction from cadmium).

3. **BISMUTH ION (Bi<sup>3+</sup>) :**

● **Precipitation with H<sub>2</sub>S in acidic medium** : Black precipitate is formed which is soluble in cold dilute HNO<sub>3</sub> and ammonium sulphide.




● **Alkaline sodium stannite (Sodium tetrahydroxidoantimonate (III))** : A black precipitate of metallic bismuth is obtained.



☞ The reagent must be freshly prepared and test must be carried out in cold solution.

● **Dilution with water** : Solution of bismuth salts gives white precipitate when water is added in larger quantity.



Soluble in mineral acids (dilute) but insoluble in tartaric acid (distinction from antimony) and in alkali hydroxide (distinction from tin).

● **Potassium iodide** : When the reagent is added dropwise to a solution containing Bi<sup>3+</sup> ions, a black precipitate is formed.





The precipitate dissolves in excess KI forming orange coloured soluble complex.




4. **CADMIUM ION (Cd<sup>2+</sup>) :**


● **Precipitation with H<sub>2</sub>S in acidic medium** : Yellow precipitate is formed which dissolves in hot dilute HNO<sub>3</sub>.



● **Ammonia solution (Dropwise addition)** : Ammonium hydroxide first gives white precipitate of Cd(OH)<sub>2</sub> which gets dissolve in excess of reagent forming a soluble complex.



● **Potassium cyanide** : Initially a white precipitate of Cd(CN)<sub>2</sub> is formed which in excess of reagent dissolves forming a soluble complex.



Qualitative Analysis

The colourless of CdS.



5.

**LEAD ION**

● **Precipit:** HNO<sub>3</sub>.

Pi

3

● **Dilute**

● **Pot soluti soluti**

● **P**


**IIB Group**

I pr



Qualitative Analysis


The colourless soluble complex is not too stable, therefore, reacts with H<sub>2</sub>S gas forming a yellow precipitate of CdS.




KI forms no precipitate (distinction from Copper)

5. **LEAD ION (Pb<sup>2+</sup>) :**


● **Precipitation with H<sub>2</sub>S in acidic medium :** Black precipitate is formed which is soluble in hot dilute HNO<sub>3</sub>.




● **Dilute H<sub>2</sub>SO<sub>4</sub> :** White precipitate is formed which is soluble in ammonium acetate.



● **Potassium iodide :** Yellow precipitate is formed which is soluble in excess more concentrated (6M) solution of the reagent. Yellow precipitate of PbI<sub>2</sub> is moderately soluble in boiling water to give a colourless solution.



● **Potassium dichromate :** Yellow precipitate is formed.



**II B Group (As<sup>3+</sup>, Sb<sup>3+</sup>, Sn<sup>2+</sup>, Sn<sup>4+</sup>)**

Soluble in (NH<sub>4</sub>)<sub>2</sub>S<sub>x</sub> as (NH<sub>4</sub>)<sub>3</sub>AsS<sub>4</sub>, (NH<sub>4</sub>)<sub>3</sub>SbS<sub>4</sub>, (NH<sub>4</sub>)<sub>2</sub>SnS<sub>3</sub>.

Add dilute HCl and filter.

Insoluble (As<sub>2</sub>S<sub>3</sub> + some S).  
Dissolve in conc. HNO<sub>3</sub> and divide into two parts.

I part + Ammonium molybdate.  
↓  
Yellow precipitate  
(NH<sub>4</sub>)<sub>3</sub>AsO<sub>4</sub> · 12 MoO<sub>3</sub>.

II part + NH<sub>4</sub>Cl + NH<sub>4</sub>OH + MgSO<sub>4</sub>.  
↓  
White precipitate  
(Mg(NH<sub>4</sub>)AsO<sub>4</sub>).


Soluble (SbCl<sub>3</sub> & SnCl<sub>4</sub>).  
Divide into two parts.

Add excess of water.  
↓  
White precipitate (SbOCl).  
↓  
Add tartaric acid.  
↓  
Precipitate is soluble.


II part + Fe powder or Zn dust, heat and add HgCl<sub>2</sub>.  
↓  
Black or grey precipitate (Hg).

1. **ARSENIC ION (As<sup>3+</sup>) :**

● **Precipitation with H<sub>2</sub>S in acidic medium :** Yellow precipitate is formed which is soluble in warm concentrated nitric acid, sodiumhydroxide solution and yellow ammonium sulphide.



● **Magnesia mixture :** White crystalline precipitate of magnesium ammonium arsenate Mg(NH<sub>4</sub>)AsO<sub>4</sub> · 6H<sub>2</sub>O is formed from neutral or ammonical solution. Upon treating the white precipitate with silver nitrate solution containing a few drops of acetic acid, red silver arsenate is formed (distinction from phosphate).



**Resonance** Educating for better tomorrow

Corporate Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.)-324005  
 Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in  
 Toll Free : 1800 200 2244 | 1800 258 5555 | CIN: U80302RJ2007PLC024029

MAINQUA - 13





## RESONANCE STUDY CENTRES (Self Owned)

### KOTA (Head Office):

Pre-Engineering Division: JEE (Advanced)  
Pre-Engineering Division: JEE (Main)  
Pre-Medical Division: AIIMS/ AIPMT  
Tel.: 0744-3012222, 3192222, 6635555  
e-mail: contact@resonance.ac.in

**Commerce & Law Program Division (CLPD)**  
Tel.: 0744-3192229, 6060663  
e-mail: clpd@resonance.ac.in

**PCCP**  
**J-2 Campus:** Tel.: 0744-2434727, 8824078330  
e-mail: pccp@resonance.ac.in

**Station Campus:** Tel.: 0744-3192223, 2440488  
e-mail: pccp.stn@resonance.ac.in

**PSPD/ MEX Division:** Tel.: 0744-2422030  
e-mail: pspd@resonance.ac.in, mex@resonance.ac.in

**DLPD**  
Tel.: 0744-6635556, 3012222  
e-mail: dlpd@resonance.ac.in

**eLPD**  
Tel.: 0744-3058242  
e-mail: elpd@resonance.ac.in

**JAIPUR**  
Tel.: 0141-8060661/ 64, 3103666, 6060662/63  
e-mail: jaipur@resonance.ac.in

**BHOPAL**  
Tel.: 0755-3206353, 3192222  
e-mail: bhopal@resonance.ac.in

**NEW DELHI**  
Tel.: 011-6060660/ 2/ 3/ 4/ 5/ 6/ 7  
e-mail: delhi@resonance.ac.in

**LUCKNOW**  
Tel.: 0522-3192222, 3192223, 6060660/ 61/ 62  
e-mail: lko@resonance.ac.in

**KOLKATA**  
Tel.: 033-31922222, 60606600/ 01/ 02  
email: kolkata@resonance.ac.in

**NAGPUR**  
Tel.: 0712-3192222, 6060660  
e-mail: nagpur@resonance.ac.in

**NANDED**  
Tel.: 02462-250220  
e-mail: nanded@resonance.ac.in

**MUMBAI**  
Tel.: 022-31922222, 60606600  
e-mail: andheri@resonance.ac.in

**UDAIPUR**  
Tel.: 0294-6060660, 5107858, 6060662  
e-mail: udaipur@resonance.ac.in

**BHUBANESWAR**  
Tel.: 0674-3192222, 3274919, 6060660/ 61  
e-mail: bbsr@resonance.ac.in

**AHMEDABAD**  
Tel.: 079-3192222/ 3/ 4 & 079-60606600/ 1/ 2  
e-mail: abed@resonance.ac.in

**PATNA**  
Tel.: 0612-3192222/ 3  
e-mail: patna@resonance.ac.in

**JODHPUR**  
Tel.: 0291- 6060660  
e-mail: jodhpur@resonance.ac.in

**AJMER**  
Tel.: 0145-3192222, 6060660/ 65  
e-mail: ajmer@resonance.ac.in

**INDORE**  
Tel.: 0731-4039100, 6060660  
e-mail: indore@resonance.ac.in

**SIKAR**  
Tel.: 01572-319222, 6060666  
e-mail: sikar@resonance.ac.in

**AGRA**  
Tel.: 0562-3192222  
e-mail: agra@resonance.ac.in

**RANCHI**  
Tel.: 0651-6060660  
e-mail: ranchi@resonance.ac.in

**ALLAHABAD**  
Tel.: 0532-6060660  
e-mail: allahabad@resonance.ac.in

**NASHIK**  
Tel.: 0253-6090028  
e-mail: nashik@resonance.ac.in

**RAIPUR**  
Tel.: 0771- 6060660  
e-mail: raipur@resonance.ac.in

**AURANGABAD**  
Tel.: 0240-6060660  
e-mail: aurangabad@resonance.ac.in

**JABALPUR**  
Tel.: 0761- 6060660  
e-mail: jabalpur@resonance.ac.in

**GWALIOR**  
Tel.: 0751-6060660  
e-mail: gwalior@resonance.ac.in

**CHANDRAPUR**  
Tel.: 07172-6060666  
e-mail: chandrapur@resonance.ac.in

**SURAT**  
Tel.: 0261-6060660  
e-mail: surat@resonance.ac.in

**RAJKOT**  
Tel.: 0281-6002011  
e-mail: rajkot@resonance.ac.in

**VADODARA**  
Tel.: 0265-6060660  
e-mail: vadodara@resonance.ac.in

## BASE STUDY CENTRES

**Base Education Service Pvt. Ltd.**  
**Bengaluru (Main Branch):**  
**Reg. Office :** No.27, Next to Indian Oil  
Petrol Bunk, Bull Temple Road,  
Basavanagudi, Bengaluru- 560004  
Tel. No.: 42604600/ 95381 41504  
**E-Mail :** info@base-edu.in  
**Website:** www.base-edu.in

**BANASANKARI II STAGE**  
Tel.: 26710835/26710836

**BELAGAVI**  
Tel.: 0831- 4208687 | Mobile: 9845226000

**CHITRADURGA**  
Mobile: 9886464755, 9972413844

**HUBLI**  
Tel.: 0836- 2252685 | Mobile: 9844118615

**INDIRANAGAR**  
Tel.: 41179342/25201306

**KALYAN NAGAR**  
Tel.: 080-25443363/25443364

**KORAMANGALA**  
Tel.: 40925512/40925534

**MALLESHWARAM**  
Tel.: 41400008

**MYSURU**  
Tel.: 0821-4242100 / 4258100/4243100

**RAJAJINAGAR**  
Tel.: 08023327588/41162135

**SHIVAMOGGA**  
Tel.: 08182-223980, 8884849590

**TUMAKURU**  
Tel.: 0816-2252387

**UDUPI**  
Tel.: 0820-2522449, 2522994, 9986663074

**VIJAYANAGAR**  
Tel.: 23111333/23111334

**YELAHANKA**  
Tel.: 08028463922/42289643

**CHIKKAMAGALURU**  
Mobile: 7411329369, 9448396890

**HASSAN**  
Mobile: 9481392014, 9972038283

**J P NAGAR**  
Tel.: 26595151/26595153.

**KALABURGI**  
Tel.: 08472-230914  
Mobile: 9845905200/9844510914



**Resonance**<sup>®</sup>  
Educating for better tomorrow

**Corporate Office:** CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Rajasthan)- 324005  
**Reg. Office:** J-2, Jawahar Nagar Main Road, Kota (Raj.)- 05 | **Tel. No.:** 0744-3192222, 3012222, 6635555 | **CIN:** U80302RJ2007PLC024029  
**To Know more:** sms **RESO** at **56677** | **E-mail:** contact@resonance.ac.in | **Website:** www.resonance.ac.in